

A LOWER BOUND ON THE LYAPUNOV EXPONENT FOR THE GENERALIZED HARPER'S MODEL

SVETLANA JITOMIRSKAYA AND WENCAI LIU

ABSTRACT. We obtain a lower bound for the Lyapunov exponent of a family of discrete Schrödinger operators $(Hu)_n = u_{n+1} + u_{n-1} + 2a_1 \cos 2\pi(\theta + n\alpha)u_n + 2a_2 \cos 4\pi(\theta + n\alpha)u_n$, that incorporates both a_1 and a_2 , thus going beyond the Herman's bound.

Dedicated to David Ruelle and Yasha Sinai on the occasion of their 80th birthdays

1. INTRODUCTION

While Lyapunov exponents of analytic one-frequency Schrödinger cocycles are positive when the coupling constants are large (uniformly in frequency and in energy)[3, 12], in general, there have been no explicit/quantitative lower estimates. The only such known explicit estimates are celebrated Herman's subharmonicity bounds: for $v(x) = \sum_{j=-M}^M a_j e^{2\pi j x}$, $a_j = \bar{a}_{-j}$, there is a uniform bound: $L(E) \geq \ln |a_M|$. This is of course not effective if $|a_M|$ is small. In this paper we present the first explicit bound that takes into account other coefficients.

Namely, we consider the discrete Schrödinger operator of the following form:

$$(1) \quad (Hu)_n = u_{n+1} + u_{n-1} + 2a_1 \cos 2\pi(x + n\alpha)u_n + 2a_2 \cos 4\pi(x + n\alpha)u_n,$$

where $a_1, a_2 \in \mathbb{R}$. It is known in physics literature as generalized Harper's model (e.g. [7, 13]), of special interest because of its connection to the three dimensional quantum Hall effect[7, 11].

Analyzing the solution of the equation $Hu = Eu$ from the dynamical systems point of view leads to the consideration of Schrödinger cocycle (α, A) , where

$$(2) \quad A(x) = \begin{pmatrix} E - 2a_1 \cos 2\pi x - 2a_2 \cos 4\pi x & -1 \\ 1 & 0 \end{pmatrix}.$$

The Lyapunov exponent (LE) of (α, A) is given by

$$L^E = \lim_{n \rightarrow \infty} \frac{1}{n} \int_{\mathbb{R}/\mathbb{Z}} \ln \|A_n(x)\| dx,$$

where

$$A_n(x) = A(x + (n-1)\alpha)A(x + (n-2)\alpha) \cdots A(x) \text{ for } n \geq 0$$

and $A_{-n}(x) = A_n(x - n\alpha)^{-1}$. We call $A_n(x)$ the n step transfer matrix. Notice that A depends on E ; sometimes we will drop this dependence from the notations, for simplicity.

By Herman's arguments [6], we have that $L^E \geq \ln |a_2|$ for all $E \in \mathbb{R}$. This yields that $L^E > 0$ if $|a_2| > 1$. Also, $L^E \geq \ln |a_1|$ if $a_2 = 0$. By the continuity of Lyapunov exponent in the cocycle (Theorem 2.1), we still have $L^E > 0$ if $|a_1| > 1$ and a_2 is small. However, Herman's method does not work to obtain a lower bound no matter how small a nonzero a_2 is. In [3, 12] a lower bound is obtained abstractly, for any analytic v in terms of its analytic extension, but in both cases in ways that so far have not led to concrete analytic bounds. In this paper we obtain a quantitative lower bound on the Lyapunov exponent, in terms of $\ln |a_1|$ and $|\frac{a_2}{a_1}|^{\frac{1}{2}}$.

Theorem 1.1. *Suppose $|a_1| > 1$ and $|a_2| < \frac{1}{100}|a_1|$. Then for all $E \in \mathbb{R}$,*

$$(3) \quad L^E \geq \ln |a_1| - 10 \left| \frac{a_2}{a_1} \right|^{\frac{1}{2}}.$$

Our result is obtained by combining the basics of Avila's global theory [1] with Herman's subharmonicity method [6] as developed by Bourgain [3], coupled with some elementary geometric considerations.

Remark. After our result was obtained, we learned that Marx, Shou, and Wellens obtained a general lower bound for the Lyapunov exponent improving Herman's results, also using the global theory [10]. However it is difficult to extract analytic quantitative results for a concrete potential from their estimates, although numerical results are possible, leading to estimates better than ours for certain parameters.

2. AVILA'S GLOBAL THEORY

Denote by $C^\omega(\mathbb{T}, \mathrm{SL}(2, C))$ the class of 1-periodic functions on $\mathbb{C}$, with analytic extension to some strip $|\Im z| < \delta$, attaining values in $\mathrm{SL}(2, C)$. Notice that matrix A given by (2) is in $C^\omega(\mathbb{T}, \mathrm{SL}(2, C))$. The following theorem shows the continuity of LE in the analytic category.

Theorem 2.1. *([4, 8]) The Lyapunov exponent $L(\alpha, D)$ is a continuous function in $\mathbb{R} \times C^\omega(\mathbb{T}, \mathrm{SL}(2, C))$ at every (α, A) with $\alpha \in \mathbb{R} \setminus \mathbb{Q}$.*

The global theory is based on the complexification of the cocycle. More precisely, given any analytic cocycle (α, D) , we consider its analytic extension, $(\alpha, D(x + i\epsilon))$ with $|\epsilon| < \delta$ for some appropriate $\delta > 0$. $L(\alpha, D_\epsilon)$ is referred to the Lyapunov exponent of the complexified cocycle $(\alpha, D(x + i\epsilon))$. Let $L(\alpha, D) = L(\alpha, D_0)$.

The Lyapunov exponent $L(\alpha, D_\epsilon)$ is known to be a convex function of ϵ (e.g. Proposition A.1, [9]). Thus we can introduce the acceleration of (α, D) ,

$$\omega(\alpha, D; \epsilon) = \frac{1}{2\pi} \lim_{h \rightarrow 0+} \frac{L(\alpha, D_{\epsilon+h}) - L(\alpha, D_\epsilon)}{h}.$$

It follows from convexity and continuity of the Lyapunov exponent that the acceleration is an upper semi-continuous function with respect to parameter ϵ [9].

The acceleration has an important property, which is stated below.

Theorem 2.2. *(Quantization of acceleration [1, 2]) Suppose $D \in C^\omega(\mathbb{T}, \mathrm{SL}(2, C))$ and $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, then we have $\omega(\alpha, D; \epsilon) \in \mathbb{Z}$.*

Applying Theorem 2.2 to Schrödinger cocycle (α, A) , where A is given by (2), we can obtain the following lemma. Below, we always assume A is given by (2), $L^E(\epsilon) = L(\alpha, A(\cdot + i\epsilon))$ and $L^E = L^E(0)$. By the symmetry ($L^E(\epsilon) = L^E(-\epsilon)$) and convexity of $L^E(\epsilon)$, we know that $\omega(\alpha, D; 0) \geq 0$.

Lemma 2.3. *For $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, there exists an $\epsilon' > 0$ such that $L^E(\epsilon) = 4\pi\epsilon + \ln |a_2|$ for $\epsilon > \epsilon'$.*

Proof. First, uniformly for $x \in \mathbb{T} = \mathbb{R}/\mathbb{Z}$, one has

$$A(x + i\epsilon) = a_2 e^{4\pi\epsilon} e^{-4\pi i x} (A_\infty + o(1)),$$

as $\epsilon \rightarrow +\infty$, where

$$A_\infty = \begin{pmatrix} -1 & 0 \\ 0 & 0 \end{pmatrix}.$$

By the continuity of the LE,

$$L^E(\epsilon) = 4\pi\epsilon + \ln |a_2|,$$

as $\epsilon \rightarrow +\infty$.

By the quantization of acceleration, we obtain

$$L^E(\epsilon) = 4\pi\epsilon + \ln|a_2|, \text{ for all } \epsilon > 0 \text{ sufficiently large.}$$

□

In addition to the convexity and continuity of L^E with respect to ϵ , we have that the graph $(\epsilon, L^E(\epsilon))$ must be one of the four pictures below (we assume here $L^E > 0$).




Fig.1




Fig.2




Fig.3




Fig.4

$\ell_2 : L^E(\epsilon) = 4\pi\epsilon + \ln|a_2|$, $\ell_1 : L^E(\epsilon) = 2\pi\epsilon + L^E$, $\ell'_1 : L^E(\epsilon) = 2\pi\epsilon + C$, $\ell_3 : L^E(\epsilon) = L^E$.

We only present the graph for $L^E > 0$, for $L^E = 0$, it is similar.

Remark 2.4. Note that the graph $(\epsilon, L^E(\epsilon))$ depends on E . The graph can determine the spectrum $\sigma(H)$.

- $\{E \in \mathbb{R} : L^E = 0\} \subset \sigma(H)$.
- If $L^E > 0$, then $E \in \sigma(H)$ if and only if the graph of $L^E(\epsilon)$ is either Fig.1 or Fig.2[1].

3. THE PROOF OF THEOREM 1.1

Before giving the proof of Theorem 1.1, two lemmas are necessary.

Lemma 3.1. Suppose $B_j = \begin{pmatrix} v_j & -1 \\ 1 & 0 \end{pmatrix}$, and $|v_j| > 2$, then we have

$$\|\prod_{j=1}^n B_j\| \geq \prod_{j=1}^n (|v_j| - 1).$$

Proof. Lemma 3.1 can be proved by induction. For this purpose, let

$$\prod_{j=1}^k B_j = \begin{pmatrix} b_{11}^k & b_{12}^k \\ b_{21}^k & b_{22}^k \end{pmatrix}.$$

Actually, we can prove the following two inequalities,

$$(4) \quad |b_{11}^k| \geq \prod_{j=1}^k (|v_j| - 1)$$

and

$$(5) \quad |b_{21}^k| \leq |b_{11}^k|.$$

Clearly, (4) and (5) hold for $k = 1$. Suppose (4) and (5) hold for k , then by induction,

$$\begin{aligned} |b_{11}^{k+1}| &= |v_{k+1}b_{11}^k - b_{21}^k| \\ &\geq |v_{k+1}||b_{11}^k| - |b_{21}^k| \\ &\geq \prod_{j=1}^{k+1} (|v_j| - 1), \end{aligned}$$

and

$$|b_{21}^{k+1}| = |b_{11}^k| \leq |b_{11}^{k+1}|.$$

This implies (4) and (5) hold for $k + 1$. □

Lemma 3.2. Suppose $e^{4\pi\epsilon_0} \geq 10$ and $a_1 > 0$. Then for any $E \in \mathbb{R}$, we have

$$\sup_{\delta \in \{\epsilon_0, 2\epsilon_0\}} \inf_{x \in \mathbb{R}} |E - a_1 e^{-2\pi\delta} e^{2\pi i x} - a_1 e^{2\pi\delta} e^{-2\pi i x}| \geq \frac{19}{60} a_1 e^{4\pi\epsilon_0}.$$

Proof. Let $a_\delta = a_1 e^{2\pi\delta} + a_1 e^{-2\pi\delta}$ and $b_\delta = a_1 e^{2\pi\delta} - a_1 e^{-2\pi\delta}$ with $\delta \in \{\epsilon_0, 2\epsilon_0\}$. Then it is easy to check that

$$\inf_{x \in \mathbb{R}} |E - a_1 e^{-2\pi\delta} e^{2\pi i x} - a_1 e^{2\pi\delta} e^{-2\pi i x}|$$

is equal to the distance $\text{dist}(E, S_\delta)$ between point $(E, 0)$ and ellipse S_δ given by

$$\frac{x^2}{a_\delta^2} + \frac{y^2}{b_\delta^2} = 1.$$

If $|E| \geq \frac{a_{\epsilon_0} + a_{2\epsilon_0}}{2}$, one has

$$\begin{aligned} \text{dist}(E, S_{\epsilon_0}) &= \inf_{x \in \mathbb{R}} |E - a_1 e^{-2\pi\epsilon_0} e^{2\pi i x} - a_1 e^{2\pi\epsilon_0} e^{-2\pi i x}| \\ &= |E - a_1 e^{-2\pi\epsilon_0} - a_1 e^{2\pi\epsilon_0}| \\ &\geq \frac{19}{60} a_1 e^{4\pi\epsilon_0}. \end{aligned}$$

If $\frac{a_{2\epsilon_0}^2 - b_{2\epsilon_0}^2}{a_{2\epsilon_0}} \leq |E| \leq \frac{a_{\epsilon_0} + a_{2\epsilon_0}}{2}$, we have

$$\begin{aligned} \text{dist}(E, S_{2\epsilon_0}) &= \inf_{x \in \mathbb{R}} |E - a_1 e^{-4\pi\epsilon_0} e^{2\pi i x} - a_1 e^{4\pi\epsilon_0} e^{-2\pi i x}| \\ &= |E - a_1 e^{-4\pi\epsilon_0} - a_1 e^{4\pi\epsilon_0}| \\ &\geq \frac{19}{60} a_1 e^{4\pi\epsilon_0}. \end{aligned}$$

If $|E| \leq \frac{a_{2\epsilon_0}^2 - b_{2\epsilon_0}^2}{a_{2\epsilon_0}}$, we also have

$$\begin{aligned} \text{dist}(E, S_{2\epsilon_0}) &= |E - a_1 e^{-4\pi\epsilon_0} e^{2\pi i x_E} - a_1 e^{4\pi\epsilon_0} e^{-2\pi i x_E}| \\ &\geq \frac{19}{60} a_1 e^{4\pi\epsilon_0}, \end{aligned}$$

where $\cos 2\pi x_E = \frac{a_{2\epsilon_0} E}{a_{2\epsilon_0}^2 - b_{2\epsilon_0}^2}$.

Putting all the cases together, we get

$$\sup_{\delta \in \{\epsilon_0, 2\epsilon_0\}} \inf_{x \in \mathbb{R}} |E - a_1 e^{-2\pi\delta} e^{2\pi i x} - a_1 e^{2\pi\delta} e^{-2\pi i x}| \geq \frac{19}{60} a_1 e^{4\pi\epsilon_0}.$$

□

Proof of Theorem 1.1

Proof. Since the Lyapunov exponent $L(\alpha, A)$ is upper semi-continuous with respect to α [5], we only need to prove Theorem 1.1 for $\alpha \in \mathbb{R} \setminus \mathbb{Q}$. Without loss of generality, assume $a_1, a_2 > 0$. Let $\epsilon_0 > 0$ be such that $e^{4\pi\epsilon_0} = (\frac{a_1}{a_2})^{\frac{1}{2}} \geq 10$. Applying Lemma 3.2, we have for any $E \in \mathbb{R}$, there exists $\delta \in \{\epsilon_0, 2\epsilon_0\}$ such that

$$(6) \quad \inf_{x \in \mathbb{R}} |E - a_1 e^{-2\pi\delta} e^{2\pi i x} - a_1 e^{2\pi\delta} e^{-2\pi i x}| \geq \frac{19}{60} a_1 e^{4\pi\epsilon_0}.$$

Case 1 : $\delta = \epsilon_0$.

We first estimate the left upper element of matrix A . Computing directly, one has for any $E, x \in \mathbb{R}$,

$$\begin{aligned} &|E - 2a_1 \cos 2\pi(x + i\epsilon_0) - 2a_2 \cos 4\pi(x + i\epsilon_0)| \\ &= |E - a_1 e^{2\pi\epsilon_0} e^{-2\pi i x} - a_1 e^{-2\pi\epsilon_0} e^{2\pi i x} - a_2 e^{4\pi\epsilon_0} e^{-4\pi i x} - a_2 e^{-4\pi\epsilon_0} e^{4\pi i x}| \\ &\geq \frac{19}{60} a_1 e^{4\pi\epsilon_0} - a_2 e^{4\pi\epsilon_0} - a_2 e^{-4\pi\epsilon_0} \\ &\geq a_1 e^{4\pi\epsilon_0} \left(\frac{19}{60} - \frac{a_2}{a_1} - \frac{a_2}{a_1} e^{-8\pi\epsilon_0} \right) \\ &> 2. \end{aligned}$$

By Lemma 3.1, we have that the left upper element of $A_n(x + i\epsilon_0)$ satisfies the following estimate

$$\begin{aligned} &|(A_n(x + i\epsilon_0) \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix})| \\ &\geq \prod_{j=0}^{n-1} (|E - 2a_1 \cos 2\pi(x + i\epsilon_0 + j\alpha) - 2a_2 \cos 4\pi(x + i\epsilon_0 + j\alpha)| - 1). \end{aligned}$$

This implies

$$(7) \quad \|A_n(x + i\epsilon_0)\| \geq \prod_{j=0}^{n-1} (|E - 2a_1 \cos 2\pi(x + i\epsilon_0 + j\alpha) - 2a_2 \cos 4\pi(x + i\epsilon_0 + j\alpha)| - 1).$$

Furthermore, we have

$$\begin{aligned}
L^E(\epsilon_0) &= \lim_{n \rightarrow \infty} \frac{1}{n} \int_{\mathbb{T}} \ln |A_n(x + i\epsilon_0)| dx \\
&\geq \int_{\mathbb{T}} \ln (|E - 2a_1 \cos 2\pi(x + i\epsilon_0) - 2a_2 \cos 4\pi(x + i\epsilon_0)| - 1) dx \\
&\geq \int_{\mathbb{T}} \ln |E - 2a_1 \cos 2\pi(x + i\epsilon_0)| dx + \int_{\mathbb{T}} \ln \left(1 - \frac{|2a_2 \cos 4\pi(x + i\epsilon_0)| + 1}{|E - 2a_1 \cos 2\pi(x + i\epsilon_0)|}\right) dx \\
&= I + II.
\end{aligned}$$

Now we will estimate I and II separately. First, one has

$$\begin{aligned}
II &\geq \ln \left(1 - \frac{60(a_2 e^{4\pi\epsilon_0} + a_2 e^{-4\pi\epsilon_0} + 1)}{a_1 e^{4\pi\epsilon_0}}\right) \\
&\geq -10 \left(\frac{a_2}{a_1}\right)^{\frac{1}{2}},
\end{aligned}$$

where the first inequality holds by (6).

We use Herman's subharmonic method to estimate I:

$$\begin{aligned}
I &= \int_{\mathbb{T}} \ln |E - a_1 e^{2\pi\epsilon_0} e^{-2\pi ix} - a_1 e^{-2\pi\epsilon_0} e^{2\pi ix}| dx \\
&= \int_{|z|=1} \ln |Ez - a_1 e^{2\pi\epsilon_0} - a_1 e^{-2\pi\epsilon_0} z^2| dz \\
&\geq 2\pi\epsilon_0 + \ln a_1.
\end{aligned}$$

Thus

$$L^E(\epsilon_0) \geq 2\pi\epsilon_0 + \ln a_1 - 10 \left(\frac{a_2}{a_1}\right)^{\frac{1}{2}}.$$

Suppose point $(\epsilon_0, L^E(\epsilon_0)) \in \ell_2$. Then one has

$$\ln a_2 + 4\pi\epsilon_0 \geq 2\pi\epsilon_0 + \ln a_1 - 10 \left(\frac{a_2}{a_1}\right)^{\frac{1}{2}}.$$

This implies that

$$\frac{40}{3} \left(\frac{a_2}{a_1}\right)^{\frac{1}{2}} \geq \ln \frac{a_1}{a_2}.$$

However this is impossible because $a_2 \leq \frac{a_1}{100}$.

Now that $(\epsilon_0, L^E(\epsilon_0)) \notin \ell_2$, we must have

$$\begin{aligned}
L^E + 2\pi\epsilon_0 &\geq L^E(\epsilon_0) \\
&\geq 2\pi\epsilon_0 + \ln a_1 - 10 \left(\frac{a_2}{a_1}\right)^{\frac{1}{2}},
\end{aligned}$$

This implies

$$L^E \geq \ln a_1 - 10 \left(\frac{a_2}{a_1}\right)^{\frac{1}{2}}.$$

Case 2: $\delta = 2\epsilon_0$.

In this case, we have

$$\inf_{x \in \mathbb{R}} |E - a_1 e^{-4\pi\epsilon_0} e^{2\pi ix} - a_1 e^{4\pi\epsilon_0} e^{-2\pi ix}| \geq \frac{19}{60} a_1 e^{4\pi\epsilon_0}.$$

The proof of case 2 is similar to that of case 1. We give the details below. Computing directly, one has for any $E, x \in \mathbb{R}$,

$$|E - 2a_1 \cos 2\pi(x + 2i\epsilon_0) - 2a_2 \cos 4\pi(x + 2i\epsilon_0)|$$

$$\begin{aligned}
&= |E - a_1 e^{4\pi\epsilon_0} e^{-2\pi ix} - a_1 e^{-4\pi\epsilon_0} e^{2\pi ix} - a_2 e^{\pi\epsilon_0} e^{-4\pi ix} - a_2 e^{-8\pi\epsilon_0} e^{4\pi ix}| \\
&\geq \frac{19}{60} a_1 e^{4\pi\epsilon_0} - a_2 e^{8\pi\epsilon_0} - a_2 e^{-8\pi\epsilon_0} \\
&\geq a_1 e^{4\pi\epsilon_0} \left(\frac{19}{60} - e^{4\pi\epsilon_0} \frac{a_2}{a_1} - e^{-12\pi\epsilon_0} \frac{a_2}{a_1} \right) \\
&> 2.
\end{aligned}$$

As in (7), we have

$$||A_n(x + 2i\epsilon_0)|| \geq \prod_{j=0}^{n-1} (|E - 2a_1 \cos 2\pi(x + 2i\epsilon_0 + j\alpha) - 2a_2 \cos 4\pi(x + 2i\epsilon_0 + j\alpha)| - 1).$$

Then we have

$$\begin{aligned}
L^E(2\epsilon_0) &= \lim_{n \rightarrow \infty} \frac{1}{n} \int_{\mathbb{T}} \ln ||A_n(x + 2i\epsilon_0)|| dx \\
&\geq \int_{\mathbb{T}} \ln (|E - 2a_1 \cos 2\pi(x + 2i\epsilon_0) - 2a_2 \cos 4\pi(x + 2i\epsilon_0)| - 1) dx \\
&\geq \int_{\mathbb{T}} \ln |E - 2a_1 \cos 2\pi(x + 2i\epsilon_0)| dx + \int_{\mathbb{T}} \ln \left(1 - \frac{|2a_2 \cos 4\pi(x + 2i\epsilon_0)| + 1}{|E - 2a_1 \cos 2\pi(x + 2i\epsilon_0)|} \right) dx \\
&= I + II.
\end{aligned}$$

Following the discussion of case 1, I and II have the following lower bounds:

$$\begin{aligned}
II &\geq \ln \left(1 - \frac{60}{19} \frac{(a_2 e^{8\pi\epsilon_0} + a_2 e^{-8\pi\epsilon_0} + 1)}{a_1 e^{4\pi\epsilon_0}} \right) \\
&\geq -10 \left(\frac{a_2}{a_1} \right)^{\frac{1}{2}}.
\end{aligned}$$

and

$$\begin{aligned}
I &= \int_{\mathbb{T}} \ln |E - a_1 e^{4\pi\epsilon_0} e^{-2\pi ix} - a_1 e^{-4\pi\epsilon_0} e^{2\pi ix}| dx \\
&\geq 4\pi\epsilon_0 + \ln a_1.
\end{aligned}$$

Thus

$$L^E(2\epsilon_0) \geq 4\pi\epsilon_0 + \ln a_1 - 10 \left(\frac{a_2}{a_1} \right)^{\frac{1}{2}}.$$

Suppose point $(2\epsilon_0, L^E(2\epsilon_0)) \in \ell_2$. Then one has

$$\ln a_2 + 8\pi\epsilon_0 \geq 4\pi\epsilon_0 + \ln a_1 - 10 \left(\frac{a_2}{a_1} \right)^{\frac{1}{2}}.$$

This implies that

$$20 \left(\frac{a_2}{a_1} \right)^{\frac{1}{2}} \geq \ln \frac{a_1}{a_2},$$

impossible because $a_2 \leq \frac{a_1}{100}$.

Now that $(2\epsilon_0, L^E(2\epsilon_0)) \notin \ell_2$, we must have

$$\begin{aligned}
L^E + 4\pi\epsilon_0 &\geq L^E(2\epsilon_0) \\
&\geq 4\pi\epsilon_0 + \ln a_1 - 10 \left(\frac{a_2}{a_1} \right)^{\frac{1}{2}}.
\end{aligned}$$

therefore

$$L^E \geq \ln a_1 - 10 \left(\frac{a_2}{a_1} \right)^{\frac{1}{2}}.$$

□

ACKNOWLEDGMENTS

S.J. is a 2014-15 Simons Fellow. This research was partially supported by NSF DMS-1401204.

REFERENCES

- [1] A. Avila. Global theory of one-frequency Schrödinger operators. *Acta Math.*, 215(1):1–54, 2015.
- [2] A. Avila, S. Jitomirskaya, and C. Sadel. Complex one-frequency cocycles. *J. Eur. Math. Soc. (JEMS)*, 16(9):1915–1935, 2014.
- [3] J. Bourgain. *Green’s Function Estimates for Lattice Schrödinger Operators and Applications.(AM-158)*. Number 158. Princeton University Press, 2005.
- [4] J. Bourgain and S. Jitomirskaya. Continuity of the Lyapunov exponent for quasiperiodic operators with analytic potential. *J. Statist. Phys.*, 108(5-6):1203–1218, 2002. Dedicated to David Ruelle and Yasha Sinai on the occasion of their 65th birthdays.
- [5] W. Craig and B. Simon. Subharmonicity of the Lyapunov index. *Duke Math. J.*, 50(551-560):132, 1983.
- [6] M. R. Herman. Une méthode pour minorer les exposants de lyapounov et quelques exemples montrant le caractere local d’un théoreme d’arnold et de moser sur le tore de dimension 2. *Commentarii Mathematici Helvetici*, 58(1):453–502, 1983.
- [7] H. Hiramoto and M. Kohmoto. Scaling analysis of quasiperiodic systems: Generalized harper model. *Physical Review B*, 40(12):8225, 1989.
- [8] S. Jitomirskaya, D. A. Koslover, and M. S. Schulteis. Localization for a family of one-dimensional quasiperiodic operators of magnetic origin. *Ann. Henri Poincaré*, 6(1):103–124, 2005.
- [9] S. Jitomirskaya and C. A. Marx. Analytic quasi-periodic cocycles with singularities and the Lyapunov exponent of extended Harper’s model. *Comm. Math. Phys.*, 316(1):237–267, 2012.
- [10] C. Marx, L. Shou, and J. Wellens. Subcritical behavior for quasi-periodic Schrödinger cocycles with trigonometric potentials. *arXiv preprint arXiv:1509.05279*, 2015.
- [11] G. Montambaux and M. Kohmoto. Quantized hall effect in three dimensions. *Physical Review B*, 41(16):11417, 1990.
- [12] E. Sorets and T. Spencer. Positive Lyapunov exponents for Schrödinger operators with quasi-periodic potentials. *Comm. Math. Phys.*, 142(3):543–566, 1991.
- [13] C. Soukoulis and E. Economou. Localization in one-dimensional lattices in the presence of incommensurate potentials. *Physical Review Letters*, 48(15):1043, 1982.

(Svetlana Jitomirskaya) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, CALIFORNIA 92697-3875, USA

E-mail address: szhitomi@math.uci.edu

(Wencai Liu) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, CALIFORNIA 92697-3875, USA

E-mail address: liuwencai1226@gmail.com